Lactacaseibacillus rhamnosus* GG and *Saccharomyces cerevisiae boulardii* supplementation exert protective effects on human gut microbiome following antibiotic administration *in vitro

C. Duysburgh¹, P. Van den Abbeele¹, M. Morera² and M. Marzorati^{1,3*}

¹ProDigest bv, Technologiepark 82, 9052 Ghent, Belgium; ²IPSEN Consumer HealthCare SAS, 65 Quai Georges Gorse, 92650 Boulogne Billancourt Cedex, France; ³Center of Microbial Ecology and Technology (CMET), Ghent University, Coupure Links 653, 9000 Ghent, Belgium; massimo.marzorati@prodigest.eu

Received: 3 September 2020 / Accepted: 7 April 2021

© 2021 Wageningen Academic Publishers

OPEN ACCESS

RESEARCH ARTICLE

Abstract

Antibiotic-induced dysbiosis of the microbial community has been associated with several gastrointestinal symptoms. The impact of repeated administration of *Lactacaseibacillus rhamnosus* GG (CNCM-I-4798) (formerly known as *Lactobacillus rhamnosus* GG), *Saccharomyces cerevisiae boulardii* (CNCM-I-1079) and their combination (associated in Smebiocta/Smectaflora Protect®) in supporting recovery of gut microbiota functionality and composition during and following amoxicillin:clavulanic acid administration was evaluated *in vitro*. Antibiotic dosage negatively affected SCFA production, coinciding with detrimental effects on *Bacteroidetes*, *Firmicutes* and *Bifidobacterium* spp. in the simulated proximal colon, while *Akkermansia muciniphila* was significantly reduced in the distal colon. *L. rhamnosus* GG and *S. boulardii* were able to thrive in both colon regions upon dosing, with *S. boulardii* even showing protective effects on the survival of *L. rhamnosus* GG during antibiotic administration. The impact of the probiotic strains on microbiome recovery revealed that supplementation with *L. rhamnosus* GG and/or *S. boulardii* resulted in a stimulating effect on the most abundant bacterial groups within the bacterial community of each donor. For one of the donors tested, co-dosing of *L. rhamnosus* GG and *S. boulardii* resulted in superior short-chain fatty acid recovery accompanied by a stronger increase in abundance of *Bifidobacteriaceae*. Overall, the current study provides first evidence that combined supplementation of *L. rhamnosus* GG and *S. boulardii* might be an interesting candidate in limiting detrimental effects of amoxicillin:clavulanic acid on the human gut microbiome, though further studies are warranted to confirm these findings.

Keywords: *Bifidobacterium*, yeast, antibiotic-associated diarrhoea, amoxicillin, clavulanic acid

1. Introduction

Although there is growing concern on aspects related to their use, antibiotics are still widely consumed. In 2017, the mean consumption of antibiotics for systemic use in Europe was 21.8 defined daily doses per 1000 inhabitants per day, with penicillins being the most commonly used class of antibiotics with a total consumption rate ranging from 36 to 71% (European Center for Disease Prevention and Control, 2018). Amoxicillin is the most widespread used penicillin-class antibiotic and is generally combined with clavulanic acid that functions as β -lactamase inhibitor,

thereby avoiding amoxicillin inactivation by β -lactamase-secreting bacteria (White *et al.*, 2004). While penicillin-class antibiotics are effective in eradicating many pathogens causing bacterial infections, they also adversely affect the indigenous gut microbial community (Brötz-Oosterhelt and Brunner, 2008). The resulting antibiotic-induced colonic imbalance has been linked with several gastrointestinal symptoms. A common disturbance includes antibiotic-associated diarrhea (AAD), which is most often caused by amoxicillin therapy (McFarland, 2008). There is a need to develop strategies that limit antibiotic-induced microbial imbalance, thereby potentially reducing

the incidence of antibiotic-associated gastrointestinal symptoms.

Probiotics are characterised as live microorganisms that confer a health benefit on the host when administered in adequate amounts, probably by supporting maintenance of a balanced gut microbiome (Hill *et al.*, 2014). While probiotic intake has been linked with a variety of physiological effects, several studies reported a potential role of probiotics as preventive and/or therapeutic agent against AAD (Agamennone *et al.*, 2018). *Lactocaseibacillus rhamnosus* GG (formerly known as *Lactobacillus rhamnosus* GG) is one of the most well-documented probiotic strains for human application and several studies reported significantly reduced prevalence of AAD upon supplementation of this probiotic strain *in vivo* (Mantegazza *et al.*, 2018; Szajewska and Kolodziej, 2015; Vanderhoof *et al.*, 1999). In fact, formulations containing a minimum dose of 2×10^9 cfu *L. rhamnosus* GG has been recommended for prevention of AAD in the Netherlands (Agamennone *et al.*, 2018). Furthermore, several other health-promoting properties have been linked with *L. rhamnosus* GG including regulation of immune function (Capurso, 2019), antipathogenic activity (De Keersmaecker *et al.*, 2006; Spacova *et al.*, 2020) and prevention of atopic disease (Kalliomaki *et al.*, 2001). Another widely used probiotic with proven health-promoting effects is the yeast *Saccharomyces cerevisiae* *boulardii* (Lazo-Vélez *et al.*, 2018). Several human studies reported significant reductions of AAD upon intake *S. boulardii* in both children and adult patients (Kotowska *et al.*, 2005; McFarland *et al.*, 1995). However, results are conflicting, with Pozzoni *et al.* (2012) and Ehrhardt *et al.* (2016) showing absence of effect of *S. boulardii* intake on the development of AAD in elderly hospitalised patients. This strengthens the need to identify specific probiotics that could accompany specific antibiotic therapies.

Despite their clinical relevance, *in vivo* studies often fail in providing mechanistic insight in how probiotics support host health. *In vitro* approaches (Minekus *et al.*, 2014, 1999; Molly *et al.*, 1993) offer a suitable alternative as strict control of environmental factors allows evaluation of cross-talk between the indigenous microbial community and the probiotic strain(s) at the site of action. However, as intestinal models depend on human faecal samples to obtain a representative microbiome, it is important to account for inter-individual variation not only during *in vivo*, but also during *in vitro* trials (Duysburgh *et al.*, 2019a).

The main objective of the current study was to evaluate the impact of repeated doses of *L. rhamnosus* GG (CNCM-I-4798), *S. boulardii* (CNCM-I-1079) and a combination of both strains in supporting recovery of gut microbiota functionality and composition upon co-supplementation with the antibiotic amoxicillin:clavulanic acid. For this purpose, the validated Simulator of the Human

Intestinal Microbial Ecosystem (SHIME®) was used (Van de Wiele *et al.*, 2015; Van den Abbeele *et al.*, 2018), thereby considering the potential inter-individual variability in microbiome composition among human individuals.

2. Materials and methods

Chemicals and test product

All chemicals were obtained from Sigma-Aldrich (Overijse, Belgium) unless stated otherwise. Lallemand Health Solutions Inc. (Blagnac, France) provided the test strains *S. boulardii* (CNCM-I-1079) and *L. rhamnosus* GG (CNCM-I-4798), associated in Smebiocta/Smectaflora Protect®. The products were administered once a day at a daily dose of 1×10^{10} *L. rhamnosus* GG (CNCM-I-4798) and/or 4×10^9 *S. boulardii* (CNCM-I-1079). The applied dosage corresponds to *in vivo* dosages as aligned with the AAD dose recommendations (Guarner *et al.*, 2017), considering the *in vitro* survival rate of the strains during upper gastrointestinal transit (results not shown).

Simulator of the Human Intestinal Microbial Ecosystem

An adaptation of the SHIME model (ProDigest and Ghent University, Ghent, Belgium) (Molly *et al.*, 1993) was used to simulate the human gastrointestinal tract. The reactor configuration was modified to a QuadSHIME setup allowing the study of four different test conditions in parallel (Dupont *et al.*, 2019). Each arm of the QuadSHIME consisted of a succession of three reactors simulating the different regions of the gastrointestinal tract, i.e. the upper gastrointestinal tract (with subsequent simulation of the gastric and small intestinal phase), the proximal colon (PC) and the distal colon (DC), respectively (Supplementary Figure S1). While the upper gastrointestinal tract reactor is of the fill-and-draw principle, the PC and DC reactors are continuously stirred and have a constant volume and pH control. The PC was operated at pH 5.6–5.9 with a retention time of 20 h, while environmental parameters in the DC included a pH-range of 6.6–6.9 and a retention time of 32 h. In order to simulate the luminal and mucus-associated microbiota, mucin-covered microcosms were included in the PC and DC (Supplementary Figure S1) as described by Van den Abbeele *et al.* (2012). Inoculum preparation, temperature settings, feeding regime and reactor feed composition were implemented from Possemiers *et al.* (2004). To evaluate the properties of four different test conditions for two healthy human adult donors (donor A: F, 24 years; donor B: F, 25 years), two parallel QuadSHIME experiments were conducted (i.e. each donor was tested in a separate QuadSHIME setup). Informed consent was retrieved for each of the donors to use the faecal samples for the study. Upon inoculation with the faecal inoculum, a two-week stabilisation period was initiated to allow the microbiome to differentiate in the different colonic reactors.

Subsequently, baseline parameters were established during a one-week control period, after which antibiotic dosage was initiated. Amoxicillin:clavulanic acid (2:1; Toku-E, Sint-Denijs-Westrem, Belgium) was administered to the PC reactors during each feeding cycle, reaching a concentration of 150 mg/day, for a period of 5 days. This antibiotic dosage was determined in order to reach the maximum *in vivo* quantity that would reach the large intestine. Indeed, *in vivo* amoxicillin is generally dosed orally at concentrations of 1,500 mg/day (De Velde *et al.*, 2016). Out of this total daily dose, 90% is absorbed in the small intestine (Legen *et al.*, 2006), resulting in a colonic dose of 150 mg/day. This colonic dose was during the current experiments dosed to the PC reactors, spread over three feeding cycles. Furthermore, *in vivo* amoxicillin is supplied in combination with clavulanic acid, with clavulanic acid being absorbed for approximately 73% (EMEA, 2001). Generally, the administered amoxicillin:clavulanic acid ratio is 4:1 or 7:1, resulting in a ratio of 1.5:1 up to 2.6:1 in the colon when considering absorption rates of 90%:73%. Therefore, an amoxicillin:clavulanic acid ratio of 2:1 was used in this study.

In the aforementioned 5-day antibiotic period, the four arms of each QuadSHIME were operated as follows: (1) control arm (CTRL), only receiving antibiotic; (2) antibiotic + *L. rhamnosus* GG (CNCM-I-4798) once per day (LGG), (3) antibiotic + *S. boulardii* (CNCM-I-1079) once per day (SB) and (4) antibiotic + a combination of *L. rhamnosus* GG (CNCM-I-4798) and *S. boulardii* (CNCM-I-1079) once per day (LGG+SB). The probiotic strains were administered once per day (i.e. during one feeding cycle) to the PC reactor, and this approximately 2 h after the antibiotic had been dosed during that feeding cycle. Administration to the PC reactor was performed as an enteric dosing strategy was envisioned.

After cessation of antibiotic administration, a one-week treatment period followed in which each experimental arm of QuadSHIME was further supplemented with their respective probiotic test products. Finally, a wash-out period of three weeks was initiated in which each QuadSHIME was operated as during the control period to assess the recovery of the gut microbiota.

Enumeration of probiotic strains

Starting from the control period, samples were collected three times per week from the luminal environment of all colonic reactors to determine the number of cfu of the probiotic strains by spread plating. Ten-fold dilution series were prepared from these samples in phosphate buffered saline (PBS) and subsequently transferred to petri dishes containing selective agar media. Selective quantification of *L. rhamnosus* GG was performed through plating on LAMVAB agar media (Hartemink *et al.*, 1997) and plates were subsequently incubated aerobically at 37 °C

for at least 72 h. Selective quantification of *S. boulardii* was performed through plating on Sabouraud agar media supplemented with chloramphenicol (50 mg/l) and plates were subsequently incubated aerobically at 30 °C for at least 36 h.

Functionality of gut microbiota

Starting from the control period, samples for determination of microbial functionality were collected three times per week from both the PC and DC. Measurement of short-chain fatty acid (SCFA) levels, including acetate, propionate, butyrate and total SCFA concentrations, was performed as reported by De Weirde *et al.* (2010). Lactate levels were determined using a commercially available enzymatic assay kit (R-Biopharm, Darmstadt, Germany) according to the manufacturer's instructions.

Community composition of gut microbiota

Starting from the control period, samples for microbiota profiling were collected once per week from the luminal environment in each colonic vessel. DNA was isolated as described by Boon *et al.* (2003), with some minor modifications as recently reported by Duysburgh *et al.* (2019b). Quantitative polymerase chain reaction (qPCR) assays for *Akkermansia muciniphila*, *Bacteroidetes* and *Firmicutes* phyla, *Lactobacillus* spp. and *Bifidobacterium* spp. were conducted using a QuantStudio 5 Real-Time PCR system (Applied Biosystems, Foster City, CA USA). Each sample was analysed in technical triplicate and outliers (more than 1 C_T difference) were removed. The qPCR for the *Firmicutes* and *Bacteroidetes* phyla were performed as described by Guo *et al.* (2008), while the qPCR for *A. muciniphila* was conducted as reported in Collado *et al.* (2007). The qPCR assay for *Lactobacillus* spp. was adopted from Furet *et al.* (2009), while the qPCR for *Bifidobacterium* spp. was performed as reported in Rinttilä *et al.* (2004). Subsequently, microbiota community composition was assessed by 16S-targeted Illumina sequencing analysis using isolated DNA originating from samples collected at the end of the control, treatment and wash-out periods. The genomic DNA extracts were diluted in DNase/RNase/protease-free water to obtain a concentration of 50 ng/μl, of which 30 μl was sent to LGC genomics GmbH (Germany) for library preparation and sequencing on an Illumina Miseq platform with v3 chemistry using primers 341F (5'-CCT ACG GGN GGC WGC AG-3') and 785Rmod (5'-GAC TAC HVG GGT ATC TAA KCC-3'), adopted from Klindworth *et al.* (2013).

Statistics

Comparison of normally distributed data of the different experimental periods on markers of microbial functionality and microbial community parameters was performed with

a Student's t-test for pairwise comparisons. Differences were considered significant if $P < 0.05$. For the 16S-targeted Illumina sequencing analysis, read assembly and clean-up was largely derived from the MiSeq procedure (Kozich *et al.*, 2013; Schloss and Westcott, 2011). In brief, the Mothur software (v. 1.40.5) was used to assemble reads into contigs, perform alignment-based quality filtering (alignment to the Mothur-reconstructed SILVA SEED alignment, v. 123), remove chimeras, assign taxonomy using a naïve Bayesian classifier (Wang *et al.*, 2007) and the SILVA NR software v132, and cluster contigs into operational taxonomic units (OTUs) at 97% sequence similarity. Sequences classified as *Eukaryota*, *Archaea*, chloroplasts and mitochondria and sequences that could not be classified, even not at (super) Kingdom level, were omitted. For each OTU, representative sequences were picked as the most abundant sequence within that OTU.

3. Results

Colon-region specific colonisation of microbial community

The main phyla in the microbial community of both donors under investigation included Firmicutes, *Bacteroidetes*, *Actinobacteria*, *Proteobacteria* and *Verrucomicrobia* and a minor fraction of *Synergistetes* and *Lentisphaerae*, though the latter was only detected in microbial community of donor B (Table 1). Specific colonisation patterns were observed in the different colonic regions. *Verrucomicrobia*, *Lentisphaerae* and *Synergistetes* specifically inhabited the DC, which was also observed for several families within the *Bacteroidetes* (e.g. *Rikenellaceae*) and *Firmicutes* (e.g. *Ruminococcaceae*) phylum, while *Bifidobacteriaceae* and *Veillonellaceae* primarily colonised the PC. Further, the biofilm covering the mucin-covered microcosms contained a highly diverse microbial community. A species-specific colonisation of the lumen versus the mucus layer was observed, with *Bacteroidaceae* specifically colonising the luminal environment, while higher levels of *Lachnospiraceae* and *Ruminococcaceae* were observed in the mucus layer. Further, higher abundance of *Synergistetes* was detected in the mucosal environment, while *Verrucomicrobia* were specifically present in the lumen. Finally, strong donor-dependent differences were observed in terms of microbial community composition. While donor A was mainly characterised by high proportions of *Bifidobacteriaceae*, *Lachnospiraceae* and *Akkermansiaceae* (the latter only in the DC), donor B showed high abundance of *Veillonellaceae*. Furthermore, several bacterial families could specifically be linked to one of the donors under investigation, such as *Muribaculaceae* and *Prevotellaceae* for donor A, and *Barnesiellaceae*, *Eubacteriaceae* and *Victivallaceae* for donor B.

Pervasive effects of antibiotic administration on endogenous microbiota

Antibiotic dosage in the absence of probiotic supplementation (CTRL) resulted in reduced SCFA levels in the PC of both donors tested (Figure 1). While butyrate levels were also reduced in the DC upon antibiotic administration for both donors tested, propionate production was stimulated towards the end of the antibiotic period in this colonic area. On the other hand, acetate levels were slightly reduced in the DC of donor A, while gradually increasing levels were observed during antibiotic dosage for donor B.

In terms of community composition, as assessed by qPCR (Table 2 and Supplementary Table S1), amoxicillin:clavulanic acid dosage resulted in significantly detrimental effects on *Bacteroidetes*, *Firmicutes* and *Bifidobacterium* spp. in the luminal and mucosal PC of both donors tested. *Lactobacillus* spp. levels were also strongly affected in the luminal PC of donor B, while significantly enhanced lactobacilli levels were observed in the mucosal environment as well as in the proximal lumen and mucus of donor A. In the DC, significantly reduced levels of *A. muciniphila* were observed for both donors tested. In contrast, *Bacteroidetes* levels were boosted in the DC during antibiotic administration in luminal and mucosal environment, with similar effects observed for the *Firmicutes* phylum (except in the lumen of donor B). Overall, upon cessation of antibiotic dosage, microbial activity and community composition recovered towards the end of the 28 days of experimental period.

Growth of *Lactacaseibacillus rhamnosus* GG and *Saccharomyces boulardii*

In order to monitor survivability of the administered test strains selective enumeration was performed in all reactors, thereby focussing on the metabolically active cell fraction of the probiotic strains (Figure 2). The combination of *L. rhamnosus* GG with *S. boulardii* resulted in a superior survival of *L. rhamnosus* GG during the antibiotic period in both colon regions and for both donors tested. Upon cessation of probiotic supplementation, *L. rhamnosus* GG washed out from the system, dropping below detection limit for donor A. In contrast, *S. boulardii* was not affected by antibiotic dosage and nor by *L. rhamnosus* GG presence/absence for both donors tested. Upon cessation of product administration, *S. boulardii* washed out from the system, dropping below detection limit in both colon regions and for both donors tested. For donor A, a faster wash-out was observed in the presence of *L. rhamnosus* GG.

Table 1. Colon-region specific colonisation of microbial community.¹

Phylum	Family	Donor A				Donor B			
		L		M		L		M	
		PC	DC	PC	DC	PC	DC	PC	DC
Actinobacteria	<i>Bifidobacteriaceae</i>	35.38%	19.93%	36.39%	16.44%	2.80%	2.15%	11.12%	3.50%
	<i>Coriobacteriaceae</i>	0.14%	0.38%	0.25%	1.09%	0.06%	0.12%	0.22%	4.10%
	<i>Eggerthellaceae</i>	0.00%	0.02%	0.00%	0.17%	0.00%	0.04%	0.00%	0.31%
	<i>Microbacteriaceae</i>	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.01%
Bacteroidetes	<i>Bacteroidaceae</i>	17.41%	12.66%	6.45%	12.02%	16.81%	22.96%	8.68%	23.34%
	<i>Barnesiellaceae</i>	0.00%	0.00%	0.00%	0.00%	0.00%	0.06%	0.00%	0.11%
	<i>Marinifilaceae</i>	0.00%	0.01%	0.00%	0.33%	0.00%	0.07%	0.00%	0.48%
	<i>Muribaculaceae</i>	0.00%	0.08%	0.00%	2.25%	0.00%	0.00%	0.00%	0.00%
	<i>Prevotellaceae</i>	0.00%	0.11%	0.00%	0.12%	0.00%	0.00%	0.00%	0.00%
	<i>Rikenellaceae</i>	0.03%	0.22%	0.04%	1.37%	0.00%	0.71%	0.00%	1.83%
	<i>Tannerellaceae</i>	2.04%	2.23%	2.28%	2.09%	0.95%	1.40%	3.38%	1.03%
	<i>Acidaminococcaceae</i>	2.94%	1.26%	0.47%	0.44%	0.00%	0.02%	0.00%	0.01%
Firmicutes	<i>Clostridiaceae_1</i>	0.00%	0.04%	0.01%	4.10%	0.00%	0.00%	0.00%	4.14%
	<i>Enterococcaceae</i>	0.00%	0.00%	0.03%	0.01%	0.16%	0.05%	0.06%	0.03%
	<i>Erysipelotrichaceae</i>	0.00%	0.06%	0.00%	0.32%	0.00%	0.19%	0.00%	0.46%
	<i>Eubacteriaceae</i>	0.00%	0.00%	0.00%	0.00%	0.00%	0.28%	0.00%	1.13%
	<i>Family_XIII</i>	0.00%	0.11%	0.00%	0.61%	0.00%	0.09%	0.00%	0.15%
	<i>Lachnospiraceae</i>	35.59%	31.70%	35.12%	42.51%	11.47%	19.97%	24.72%	25.98%
	<i>Lactobacillaceae</i>	0.00%	0.00%	0.00%	0.02%	0.00%	0.00%	0.00%	0.01%
	<i>Peptostreptococcaceae</i>	0.00%	0.00%	0.00%	0.09%	0.00%	0.00%	0.00%	0.00%
	<i>Ruminococcaceae</i>	0.09%	6.11%	0.80%	6.52%	0.03%	2.51%	0.30%	6.17%
	<i>Veillonellaceae</i>	0.71%	0.41%	12.61%	1.23%	67.51%	46.03%	51.10%	22.26%
Lentisphaerae	<i>Victivallaceae</i>	0.00%	0.00%	0.00%	0.00%	0.00%	0.01%	0.00%	0.11%
Proteobacteria	<i>Burkholderiaceae</i>	1.46%	0.76%	0.75%	1.95%	0.19%	0.66%	0.25%	1.49%
	<i>Desulfovibrionaceae</i>	1.97%	1.88%	3.08%	2.51%	0.00%	1.65%	0.01%	1.31%
	<i>Enterobacteriaceae</i>	0.67%	0.24%	0.37%	1.50%	0.03%	0.06%	0.07%	0.10%
	<i>Pseudomonadaceae</i>	1.00%	1.06%	0.10%	0.12%	0.00%	0.71%	0.01%	0.23%
	<i>uncultured</i>	0.54%	0.15%	1.21%	0.06%	0.00%	0.00%	0.00%	0.00%
	<i>Xanthomonadaceae</i>	0.00%	0.22%	0.00%	0.01%	0.00%	0.01%	0.00%	0.00%
Synergistetes	<i>Synergistaceae</i>	0.00%	0.02%	0.00%	0.16%	0.00%	0.03%	0.00%	1.25%
Verrucomicrobia	<i>Akkermansiaceae</i>	0.00%	20.28%	0.00%	1.42%	0.00%	0.13%	0.00%	0.14%

¹ Average abundance (%) at microbial family level in the proximal (PC) and distal colon (DC) and the luminal (L) and mucosal environment (M) of the SHIME at the end of the control period for two different human donors (n=4, corresponding to the 4 units within each SHIME) as obtained by 16S-targeted Illumina sequencing. Statistically significant differences between the PC and DC within each environment are underlined, while statistically significant differences between the luminal and mucosal environment within each colonic region are indicated in italics ($P<0.05$). The intensity of green shading indicates the absolute abundance, normalized for each of the different families per donor.

Effects of test products on the microbial metabolic activity

Recovery of microbial metabolic activity following antibiotic dosage was investigated by normalizing SCFA and lactate levels during the final week of the wash-out period to the respective control periods (Table 3). These normalised SCFA profiles resulted in treatment-specific observations.

For donor A, a significantly higher production of acetate, propionate and butyrate was observed compared to the antibiotic control incubation (CTRL) upon co-dosing *L. rhamnosus* GG and *S. boulardii* in both colon regions (except for acetate in the PC and butyrate in the DC were no significant effect, but a clear trend was observed). Furthermore, the combination of *L. rhamnosus* GG and *S. boulardii* resulted in synergetic effects as shown

Figure 1. Pervasive effect of antibiotic supplementation on metabolite production. Absolute concentrations (mM) of acetate (AA), propionate (PA), butyrate (BA) and total SCFA in the proximal (PC) and distal colon (DC) reactors for the control unit (CTRL; only receiving antibiotic and thus not supplemented with probiotics during the antibiotic and treatment period) of the SHIME for two different human donors (n=1). Samples were taken one control (C), one antibiotic (AB), one treatment (TR) and three wash-out (WO1, WO2 and WO3) weeks. During each week, three samples (A, B and C) were collected.

by significantly enhanced levels of acetate in the PC, propionate in the DC and butyrate in both colon regions when comparing the respective individual probiotic supplementation with the combined *L. rhamnosus* GG and *S. boulardii* condition. For donor B, all test conditions resulted in acetate levels that were higher compared to the control period, however no differences were observed compared to the antibiotic control incubation (CTRL). Similarly, no significant differences were observed between the different test conditions in terms of propionate production. In contrast, administration of *S. boulardii* resulted in significantly increased butyrate levels compared to the antibiotic control incubation (CTRL) in both colon regions. Finally, dosage of *L. rhamnosus* GG and its combination with *S. boulardii* significantly reduced lactate production compared to the antibiotic control (CTRL) in the DC of donor B. For donor A, no significant differences were observed in terms of lactate production.

Effects of test products on microbial community composition

To assess community composition, a quantitative assessment via qPCR was performed for targeted microbial groups (Figure 3 and Supplementary Figure S2). *L. rhamnosus* GG supplementation (with or without *S. boulardii*) strongly increased luminal and mucosal *Lactobacillus* spp. levels in both colon regions for donor A and B, which was probably a direct consequence of the addition of the *L. rhamnosus* GG strain for these test conditions, thereby indicating that high levels of the probiotic test strain could be maintained in a complex microbial environment. For donor A, levels of *Lactobacillus* spp. also increased upon amoxicillin:clavulanic acid dosage during the antibiotic control incubation (CTRL), however the observed increase was much smaller compared to the strong increase upon *L. rhamnosus* GG supplementation. During the wash-out period, *Lactobacillus* spp. levels decreased (except upon

Table 2. Pervasive effect of antibiotic administration on bacterial groups as assessed by qPCR.¹

	PC				DC			
	C	AB	TR	WO	C	AB	TR	WO
Donor A								
<i>Akkermansia muciniphila</i>	below LOD	below LOD	below LOD	below LOD	9.33±0.01	5.90±0.03	8.23±0.04	8.89±0.40
<i>Lactobacillus</i> spp.	4.43±0.29	7.13±0.04	5.32±0.09	5.63±0.64	4.69±0.13	6.61±0.04	4.79±0.08	4.75±0.35
<i>Bifidobacterium</i> spp.	9.73±0.04	8.44±0.00	9.77±0.02	10.00±0.10	9.88±0.03	9.21±0.01	9.85±0.07	9.62±0.11
<i>Bacteroidetes</i>	9.86±0.04	6.62±0.02	9.90±0.05	9.98±0.09	10.07±0.03	10.55±0.05	10.33±0.01	9.99±0.15
<i>Firmicutes</i>	9.64±0.01	8.56±0.01	9.73±0.04	9.84±0.14	9.98±2.49	10.07±4.06	9.97±4.31	9.98±3.29
Donor B								
<i>A. muciniphila</i>	below LOD	±			8.02±0.05	below LOD	6.72±0.01	8.14±3.13
<i>Lactobacillus</i> spp.	5.67±0.00	5.24±0.02	6.04±0.02	5.69±0.14	5.53±0.02	4.67±0.09	5.10±0.13	5.17±0.09
<i>Bifidobacterium</i> spp.	9.10±0.02	6.61±0.02	9.86±0.09	9.52±0.29	9.29±0.03	8.69±0.01	9.73±0.03	9.52±0.16
<i>Bacteroidetes</i>	9.88±0.01	6.76±0.03	10.56±0.05	9.76±0.21	10.26±0.02	10.65±0.02	10.47±0.05	10.22±0.11
<i>Firmicutes</i>	9.88±0.01	9.13±0.01	9.98±0.02	9.61±0.16	10.12±0.08	9.89±0.03	10.23±0.03	10.04±0.14

¹ Average *Bifidobacterium* spp., *Lactobacillus* spp., *A. muciniphila*, *Bacteroidetes* and *Firmicutes* levels (log 16S rRNA copies/ml) over the control (C; n=1 in technical triplicate), the antibiotic (AB; n=1 in technical triplicate), the treatment (TR; n=1 in technical triplicate) and the wash-out (WO; n=3 in technical triplicate) periods in the luminal proximal (PC) or distal colon (DC) for the control unit (CTRL; only receiving antibiotic) of the SHIME for two different human donors. Data is presented as mean ± standard deviation. Statistically significant differences relative to the control period are indicated in bold ($P<0.05$). The intensity of green shading indicates the absolute abundance, normalised for each of the different bacterial groups per colonic region. LOD = limit of detection.

S. boulardii administration in donor A), with no notable differences being observed between the different test conditions. Further, antibiotic dosage strongly decreased *Bifidobacterium* spp. levels for both donors tested, especially in the luminal environment. Upon cessation of antibiotic dosage, recovery of *Bifidobacterium* spp. levels was observed in all treated arms, with no notable differences between the different test conditions.

Microbiota profiling at the main site of fermentation (i.e. luminal PC) (Figure 4), using 16S-targeted Illumina sequencing, confirmed the significantly increased abundance of *Lactobacillaceae* upon supplementation of *L. rhamnosus* GG (both in the presence and absence of *S. boulardii*) for both donors tested. Similar results, though less apparent, were observed in the mucosal environment (Supplementary Figure S3). As shown by qPCR analyses, this enrichment proved to be a transient effect, as cessation of product administration resulted in a markedly reduced abundance of *Lactobacillaceae* to levels below the detection limit during the wash-out period. For donor A, antibiotic dosage resulted in reduction of the highly abundant *Bifidobacteriaceae* family, at the expense of *Bacteroidaceae*. Furthermore, *Proteobacteria* were found specifically enriched by antibiotic dosage, as seen by increased abundances of several families belonging to the *Proteobacteria* phylum for all test conditions. Following

product administration, the combination of *L. rhamnosus* GG and *S. boulardii* resulted in the strongest increase in *Bifidobacteriaceae* abundance. In the mucosal environment, the increase in *Bifidobacteriaceae* was already observed during the treatment period. Moreover, compared to the antibiotic control (CTRL), supplementation of *S. boulardii* alone or in combination with *L. rhamnosus* GG resulted in increased levels of *Lachnospiraceae*, mainly at the expense of *Veillonellaceae*. Whereas the microbial community of donor A was strongly enriched in *Bifidobacteriaceae*, the gut microbiota of donor B was characterised by high abundance of *Veillonellaceae*. Antibiotic dosage drastically reduced *Veillonellaceae* levels, as seen by the strong decrease in the antibiotic control incubation (CTRL) during the treatment period in both the luminal and mucosal environment. Remarkably, administration of the different test products alleviated this effect, as *Veillonellaceae* levels remained highly abundant under these test conditions following antibiotic dosage. Following treatment, abundance of several families within the *Proteobacteria* phylum increased for all test conditions, with a specific increase in *Enterobacteriaceae* upon *S. boulardii* supplementation in donor A. Furthermore, some product-specific effects were observed, including increased levels of *Lachnospiraceae* upon supplementation of *L. rhamnosus* GG and co-dosing *L. rhamnosus* GG with *S. boulardii* and enhanced *Bacteroidaceae* levels upon administration of *L. rhamnosus* GG.

Figure 2. Growth of *Lactocaseibacillus rhamnosus* GG (LGG) and *Saccharomyces boulardii* (SB). Average LGG and SB levels (log cfu/ml) over the control (C), the antibiotic (AB), the treatment (TR) and three wash-out (WO1, WO2 and WO3) weeks in the luminal proximal (PC) or distal colon (DC) upon administration of the individual LGG or SB strain and their combination (LGG+SB) in the SHIME for two different human donors. Data is presented as mean $\pm$ standard deviation (n=3).

4. Discussion

During the current study, the impact of repeated doses of *L. rhamnosus* GG, *S. boulardii* and a combination of both strains on recovery of gut microbiota functionality and composition upon co-administration with the antibiotic amoxicillin:clavulanic acid was investigated *in vitro*. For this purpose, the validated M-SHIME model was used, which allows to create a stable microbial community that is representative for the different colonic regions (Liu

et al., 2018; Possemiers *et al.*, 2004). The present study confirmed that the M-SHIME model resulted in the creation of a colon region-specific luminal and mucosal microbial community, representative for each of the donors under investigation. Overall, the PC was mainly enriched in bacterial families involved in primary substrate degradation, including *Bifidobacteriaceae* and *Veillonellaceae*, while the environmental conditions in the DC selected for microbial groups with specific metabolic functionalities, such as mucin degradation by *Akkermansiaceae* (Van Herreweghen

Table 3. Recovery of metabolite production.¹

	PC				DC			
	CTRL	LGG	SB	LGG+SB	C	LGG	SB	LGG+SB
Donor A								
Acetate (mM)	-2.46±4.79 ^{a,b}	-1.92±1.94 ^a	1.03±1.39 ^a	6.58±1.71 ^b	2.81±3.58 ^a	6.28±3.75 ^{a,b}	8.87±3.08 ^{a,b}	11.40±2.43 ^b
Propionate (mM)	-2.56±0.52 ^a	-2.50±0.36 ^{a,b}	-1.39±0.92 ^a	0.10±0.83 ^b	-3.18±0.24 ^a	-3.58±0.38 ^a	-3.19±1.00 ^a	0.89±0.80 ^b
Butyrate (mM)	0.25±0.72 ^a	1.50±0.61 ^b	2.33±0.18 ^b	4.75±0.26	-1.09±1.20 ^{a,b}	-2.49±0.82 ^a	-0.75±0.64 ^a	2.97±0.53 ^b
Lactate (mM)	-0.46±0.10	-0.52±0.06	-0.50±0.02	-0.53±0.02	-0.16±0.06 ^a	-0.03±0.07 ^b	0.28±0.63 ^{a,b}	1.04±0.83 ^{a,b}
Total SCFA (mM)	-0.33±4.06 ^{a,b}	0.83±2.00 ^a	3.37±1.46 ^a	12.42±2.36 ^b	2.98±2.49 ^a	6.90±4.06 ^{a,b}	11.11±4.31 ^{a,b}	16.85±3.29 ^b
Donor B								
Acetate (mM)	5.21±1.20	21.54±10.39	8.85±1.90	9.89±3.66	12.84±1.62	15.59±4.26	9.44±5.07	13.19±3.13
Propionate (mM)	-0.21±0.17	2.80±1.64	0.51±0.38	0.57±0.28	0.07±0.22	0.89±0.63	0.45±1.03	-0.03±0.41
Butyrate (mM)	-1.05±0.65 ^a	-0.94±0.56 ^a	0.64±0.81 ^b	-0.09±0.17 ^{a,b}	-0.47±0.53 ^a	-1.60±0.49 ^a	0.73±0.26 ^b	-0.92±0.31 ^a
Lactate (mM)	0.29±0.36 ^{a,b}	-0.09±0.16 ^a	0.19±0.19 ^b	0.39±0.27 ^b	-0.02±0.06 ^a	-0.26±0.03 ^b	0.01±0.18 ^{a,b}	-0.16±0.09 ^b
Total SCFA (mM)	2.62±2.03	20.39±11.85	8.85±1.51	8.72±3.49	12.48±2.02	13.33±4.91	9.56±6.07	12.20±2.94

¹ Average normalised concentration of acetate, propionate, butyrate, lactate and total short chain fatty acid (SCFA, mM) associated to the administration of *Lactocaseibacillus rhamnosus* GG (LGG), *Saccharomyces boulardii* (SB) and their combination (LGG+SB) compared to a control (CTRL) in the proximal (PC) and distal colon (DC) during the final wash-out week (WO3) in the SHIME for two different human donors. Data is presented as mean ± standard deviation (n=3). Statistically significant differences between the different conditions in each of the colonic environments are indicated with different letters ($P<0.05$). The intensity of green shading indicates the absolute abundance, normalised for each of the different metabolites per colonic region.

et al., 2017). This colon-region specificity confirms previous findings by Van den Abbeele *et al.* (2010). Further, the mucosal biofilm has been shown to be specifically colonised by butyrate-producing members of the *Firmicutes* phylum (Van den Abbeele *et al.*, 2013) as well as species from the *Synergistetes* phylum (Liu *et al.*, 2018), as demonstrated during the current study. Overall, the establishment of a representative microbiome in the M-SHIME model provided an excellent tool for performing mechanistic research to unravel effects on the human gut microbiome.

Antibiotic administration strongly affected the *in vitro* microbial community, resulting in diminished levels of SCFA and detrimental effects in terms of community composition. Overall effects were strongest in the PC, probably due to dilution of antibiotic treatment or microbial degradation of amoxicillin:clavulanic acid when moving to the DC. Butyrate production was most strongly affected, showing significant reductions in both colon regions upon amoxicillin:clavulanic acid dosage, confirming findings of *in vitro* work performed by Ladirat *et al.* (2014b). Moreover, the abundance of *Lachnospiraceae* and *Ruminococcaceae*, two families containing several butyrate-producing species, decreased significantly following amoxicillin treatment *in vivo* (Ladirat *et al.*, 2014a; Tsitko *et al.*, 2019), which was for the former bacterial family also observed during the current study in the luminal environment of both donors tested. In concordance with previous *in vivo* studies (Ladirat

et al., 2014a; Soldi *et al.*, 2019), *Bifidobacterium* spp. levels were also strongly affected by the amoxicillin:clavulanic acid administration. Propionate production was stimulated in the DC coinciding with enhanced *Bacteroidetes* proportions towards the end of the antibiotic period. While several studies have reported a similar stimulation of several members of the *Bacteroidetes* phylum following amoxicillin administration (Barc *et al.*, 2004; Panda *et al.*, 2014), others showed reduced *Bacteroidetes* levels as observed in the PC in the current study (Ladirat *et al.*, 2014b; Tsitko *et al.*, 2019). It can be hypothesised that the suppression of *Bifidobacterium* spp. and butyrate-producing microorganisms by antibiotic dosage created a competitive advantage for other bacterial species, including members of the *Bacteroidetes* phylum. This might induce enrichment of the Gram-negative bacterial flora (such as several groups within the *Bacteroidetes* phylum), which has been linked with several side effects such as the induction of diarrhoea (Ramamurthy *et al.*, 2013). Overall, it can be concluded that amoxicillin:clavulanic acid dosage exerted a detrimental effect on the *in vitro* microbial community, in accordance with *in vivo* data, which might be related with the occurrence of AAD.

In order to link the potential effect of *L. rhamnosus* GG and *S. boulardii* on the recovery of the gut microbiome during and following antibiotic supplementation, the presence of viable *L. rhamnosus* GG and *S. boulardii*

Figure 3. Luminal microbial community composition as assessed via qPCR. Average *Bifidobacterium* spp. and *Lactobacillus* spp. levels (log 16S rRNA copies/mL) associated to the administration of *Lacticaseibacillus rhamnosus* GG (LGG), *Saccharomyces boulardii* (SB) and their combination (LGG+SB) compared to a control (CTRL) in the luminal proximal (PC) or distal colon (DC) during the control (C; n=1 in technical triplicate), the antibiotic (AB; n=1 in technical triplicate), the treatment (TR; n=1 in technical triplicate) and the wash-out (WO; n=3 in technical triplicate) periods in the SHIME for two different human donors. Data is presented as mean $\pm$ standard deviation.

cells was assessed throughout the experimental period. Selective enumeration pointed out that viable *L. rhamnosus* GG cells were specifically detected in the SHIME units where *L. rhamnosus* GG was administered. The survival of *L. rhamnosus* GG during antibiotic dosage was enhanced upon co-supplementation with *S. boulardii* versus the unit where *L. rhamnosus* GG was administered alone, and this

in both colon regions and for both donors tested. Recently, it was shown by our research group that no antagonistic effects between *L. rhamnosus* GG and *S. boulardii* were observed when cultivated together (Moens *et al.*, 2019). The current study further complements these observations, showing protective effects of *S. boulardii* on survival of *L. rhamnosus* GG during antibiotic dosage. Further, high

Figure 4. Luminal microbial community composition as assessed via 16S-targeted Illumina sequencing. Abundance (%) at microbial family level associated to the administration of *Lactobacillus rhamnosus* GG (LGG), *Saccharomyces boulardii* (SB) and their combination (LGG+SB) compared to a control (CTRL) in the luminal proximal colon (PC) at the end of the control period (C), treatment period (TR) and wash-out period (WO) in the SHIME for two different human donors (n=1).

levels of viable *S. boulardii* cells were detected in the SHIME units where *S. boulardii* was administered and these levels were not affected by the antibiotic dosage, confirming literature findings (Neut *et al.*, 2017), regardless of the presence/absence of *L. rhamnosus* GG. Upon cessation of the product administration, both test strains washed out from the system. It has been shown that persistence of individual strains varies in a strain-specific manner *in vivo* (Tremblay *et al.*, 2020). Indeed, while *S. boulardii* is only to persist in the colonic environment for hours upon cessation of dosing, *L. rhamnosus* GG can persevere for several weeks in the colonic environment (McFarland and Elmer, 2005). Overall, these findings demonstrated that the strains were not only dosed but also capable to thrive in both colon regions of both donors tested, thus potentially resulting in beneficial effects on antibiotic-induced dysbiosis.

The impact of the probiotic test strains on the recovery of microbial functionality and composition showed donor-dependent differences. 16S-targeted Illumina sequencing revealed that supplementation of *L. rhamnosus* GG and/or *S. boulardii* resulted in a protective effect on the most abundant bacterial groups within the bacterial community of each donor upon antibiotic dosage. Donor A was mainly characterised by high proportions of *Bifidobacteriaceae* and *Lachnospiraceae*, with both bacterial groups being highly affected by antibiotic dosage. Upon cessation of antibiotic administration, *Lachnospiraceae* recovered in all test conditions, with higher levels observed upon probiotic supplementation, especially for addition of *S. boulardii* and its combination with *L. rhamnosus* GG. Supplementation of *S. boulardii* has been reported to lead

to a faster recovery of the gut microbiome following a state of dysbiosis, including positive effects on *Lachnospiraceae* levels (Moré and Swidsinski, 2015). Furthermore, co-dosing *L. rhamnosus* GG and *S. boulardii* resulted in superior SCFA recovery compared to the antibiotic control, accompanied by a stronger increase in abundance of *Bifidobacteriaceae*. This confirmed previous results from our research group, which demonstrated that combination of *L. rhamnosus* GG and *S. boulardii* improved microbiota functionality under impaired conditions (Moens *et al.*, 2019). Donor B instead was characterised by a higher abundance of *Veillonellaceae* in its healthy state, levels of which were drastically reduced following antibiotic dosage in the control arm, while supplementation of the different probiotic test strains preserved *Veillonellaceae* levels, indicating a potential protective effect of *L. rhamnosus* GG and *S. boulardii* on *Veillonellaceae* levels during antibiotic dosage for this particular donor. Collectively, these results indicate that effects of antibiotic co-treatment with *L. rhamnosus* GG and *S. boulardii* on gut microbiota functionality and composition may differ substantially depending on the baseline composition of the gut microbiota of each subject, although reinforcing the hypothesis that combined intake of *L. rhamnosus* GG and *S. boulardii* might stimulate microbiome recovery following antibiotic dosage.

Some potential limitations of the study are acknowledged. While inter-individual variations were evaluated, only two donors were selected for the current study. However, it is important to note that the human microbial community composition is generally characterised by large inter-individual differences (Eckburg *et al.*, 2005) and this

should therefore also be taken into account upon using *in vitro* gut models. It has been reported that the gut microbial community of healthy individuals naturally recovers following antibiotic administration, reaching a near-baseline composition within 45 days following a short antibiotic intervention period (Palleja *et al.*, 2018). In the current study, a parallel antibiotic control was included to map natural microbiome recovery following a 5-day amoxicillin:clavulanic acid dosage *in vitro*. Near-full recovery of microbial metabolic activity and community composition was observed during the post-intervention period of 28 days. This fast recovery of the microbial community during the present *in vitro* study might therefore have obscured potential effects of *L. rhamnosus*, *S. boulardii* and their combination on microbiome recovery during the post-intervention period. Furthermore, while the gut microbial community generally recovers to its healthy state in 45 days due to its resilience, long-term alterations may persist in some subjects explaining susceptibility to AAD or other dysbiosis-related diseases (De La Cochetière *et al.*, 2005; Sommer and Bäckhed, 2013). Furthermore, host-microbiome interactions were not investigated during the current study. While the current study has shown that not only the luminal microbial community was affected by the administered test products, but that also the mucus-associated microbiota was impacted, these results point to potential interesting interactions with the intestinal epithelial cells. Combining samples obtained from the SHIME model with a co-culture human cell model (Daguet *et al.*, 2016), could have provided insight on the immunomodulatory properties of *L. rhamnosus*, *S. boulardii* and their combination following antibiotic dosage.

In conclusion, administration the antibiotic amoxicillin:clavulanic acid exerted a detrimental effect on the human gut microbiome in the simulated gastrointestinal tract. While recovery of the microbial community was observed at the end of the 28d experimental period, supplementation of *L. rhamnosus* GG and *S. boulardii* positively impacted colonic balance, though in a donor-dependent fashion. Evidence was provided that co-dosing of *L. rhamnosus* GG and *S. boulardii* might additionally stimulate recovery of microbial functionality and composition following amoxicillin:clavulanic acid dosage, which could provide an interesting approach for antibiotic-related gastrointestinal symptoms in the clinical environment such as in the protection against AAD. However, future research is warranted to confirm these findings.

Supplementary material

Supplementary material can be found online at <https://doi.org/10.3920/BM2020.0180>.

Figure S1. Design of SHIME experiment.

Figure S2. Mucosal microbial community composition as assessed via qPCR.

Figure S3. Mucosal microbial community composition as assessed via 16S-targeted Illumina sequencing.

Table S1. Pervasive effect of antibiotic administration on mucosal bacterial groups as assessed by qPCR.

Acknowledgements

This work was sponsored by IPSEN Consumer HealthCare SAS. The authors thank the donors who kindly supplied a faecal sample for the purpose of the study. Further, the authors thank Dr Thomas A. Tompkins, Dr Annie Tremblay and Dr Frédéric Moens for critically reviewing the manuscript.

Conflict of interest

Mireia Morera is an employee of IPSEN Consumer HealthCare SAS.

References

- Agamennone, V., Krul, C.A.M., Rijkers, G. and Kort, R., 2018. A practical guide for probiotics applied to the case of antibiotic-associated diarrhea in the Netherlands. *BMC Gastroenterology* 18: 103.
- Barc, M.C., Bourlioux, F., Rigottier-Gois, L., Charrin-Sarnel, C., Janoir, C., Boureau, H., Dore, J. and Collignon, A., 2004. Effect of amoxicillin-clavulanic acid on human fecal flora in a gnotobiotic mouse model assessed with fluorescence hybridization using group-specific 16S rRNA probes in combination with flow cytometry. *Antimicrobial Agents and Chemotherapy* 48: 1365-1368.
- Boon, N., Top, E.M., Verstraete, W. and Siciliano, S.D., 2003. Bioaugmentation as a tool to protect the structure and function of an activated-sludge microbial community against a 3-chloroaniline shock load. *Applied and Environmental Microbiology* 69: 1511-1520.
- Brötz-Oesterhelt, H. and Brunner, N.A., 2008. How many modes of action should an antibiotic have? *Current Opinion in Pharmacology* 8: 564-573.
- Capurso, L., 2019. Thirty years of *Lactobacillus rhamnosus* GG: a review. *Journal of Clinical Gastroenterology* 53: S1-S41.
- Collado, M.C., Derrien, M., Isolauri, E., de Vos, W.M. and Salminen, S., 2007. Intestinal integrity and *Akkermansia muciniphila*, a mucin-degrading member of the intestinal microbiota present in infants, adults, and the elderly. *Applied and Environmental Microbiology* 73: 7767-7770.
- Daguet, D., Pinheiro, I., Verhelst, A., Possemiers, S. and Marzorati, M., 2016. Arabinogalactan and fructooligosaccharides improve the gut barrier function in distinct areas of the colon in the simulator of the human intestinal microbial ecosystem. *Journal of Functional Foods* 20: 369-379.

- De Keersmaecker, S.C., Verhoeven, T.L., Desair, J., Marchal, K., Vanderleyden, J. and Nagy, I., 2006. Strong antimicrobial activity of *Lactobacillus rhamnosus* GG against *Salmonella* Typhimurium is due to accumulation of lactic acid. *FEMS Microbiology Letters* 259: 89-96.
- De La Cochetière, M.F., Durand, T., Lepage, P., Bourreille, A., Galmiche, J.P. and Doré, J., 2005. Resilience of the dominant human fecal microbiota upon short-course antibiotic challenge. *Journal of Clinical Microbiology* 43: 5588-5592.
- De Velde, F., de Winter, B.C., Koch, B.C., van Gelder, T. and Mouton, J.W., 2016. Non-linear absorption pharmacokinetics of amoxicillin: consequences for dosing regimens and clinical breakpoints. *Journal of Antimicrobial Chemotherapy* 71: 2909-2917.
- De Weirtdt, R., Possemiers, S., Vermeulen, G., Moerdijk-Poortvliet, T.C.W., Boschker, H.T.S., Verstraete, W. and Van de Wiele, T., 2010. Human faecal microbiota display variable patterns of glycerol metabolism. *FEMS Microbiology Ecology* 74: 601-611.
- Dupont, D., Alric, M., Blanquet-Diot, S., Bornhorst, G., Cueva, C., Deglaire, A., Denis, S., Ferrua, M., Havenaar, R., Lelieveld, J., Mackie, A.R., Marzorati, M., Menard, O., Minekus, M., Miralles, B., Recio, I. and Van den Abbeele, P., 2019. Can dynamic *in vitro* digestion systems mimic the physiological reality? *Critical Reviews in Food Science and Nutrition* 59: 1546-1562.
- Duysburgh, C., Van Den Abbeele, P., Bothe, M., Stover, J., Kuchinka-Koch, A., Schwejda-Guettes, S. and Marzorati, M., 2019a. Lactulose crystals beneficially affect community composition along entire human colon *in vitro*, resulting in donor-dependent prebiotic effects at metabolic level. *International Journal of Nutrition and Food Sciences* 8: 10.
- Duysburgh, C., Van den Abbeele, P., Krishnan, K., Bayne, T.F. and Marzorati, M., 2019b. A synbiotic concept containing spore-forming *Bacillus* strains and a prebiotic fiber blend consistently enhanced metabolic activity by modulation of the gut microbiome *in vitro*. *International Journal of Pharmaceutics* X 1: 100021.
- Eckburg, P.B., Bik, E.M., Bernstein, C.N., Purdom, E., Dethlefsen, L., Sargent, M., Gill, S.R., Nelson, K.E. and Relman, D.A., 2005. Diversity of the human intestinal microbial flora. *Science* 308: 1635-1638.
- Ehrhardt, S., Guo, N., Hinz, R., Schoppen, S., May, J., Reiser, M., Schroeder, M.P., Schmiedel, S., Keuchel, M., Reisinger, E.C., Langeheinecke, A., de Weerth, A., Schuchmann, M., Schaberg, T., Ligges, S., Eveslage, M., Hagen, R.M., Burchard, G.D., Lohse, A.W., Group, f.t.S.S., Ruf, B., Porschen, R., Trenn, G., Butterfaß-Bahloul, T., Wuerthwein, G., Oeder, F., Runge, A., Klauss, E., Hansen-Rosenblatt, N., Werner, T., Schulze, K., Kreuels, B., Schäfer, G., Hübener, P., Hennigs, A., Beisel, C., Fischer-Brügge, D., Zimmermann-Fraedrich, K., Röder, C., Grigo, N., Riecke, A., Schreckenbauer, H., Hemmer, C., Klammt, S., Geerdes-Fenge, H., Frimmel, S., Kittner, J.M., Rey, J.W., Schattenberg, J.M., Thieringer, F., Schmits, R., Grandt, D., Büch, P.M., Klebert, A., Mittag, M.A., Lehnen, S., Tiefengraber, D., Radecke, K., Hering, I., Zeller, W., Rundt, L., Brandt, L., Baltes, P., Dajani, D., Kurniawan, N., Pflüger, C., Behjat, N., Engel, U., Unger, M. and Group, f.t.S.S., 2016. *Saccharomyces boulardii* to prevent antibiotic-associated diarrhea: a randomized, double-masked, placebo-controlled trial. *Open Forum Infectious Diseases* 3: ofw011.
- European Agency for the Evaluation of Medicinal Products (EMA), 2001. Clavulanic acid. Report EMA/MRL/776?01-Final. Committee for Veterinary Medicinal Products, London, pp. 1-6.
- European Center for Disease Prevention and Control, 2018. Antimicrobial consumption – Annual Epidemiological Report for 2017. Annual Epidemiological Report on Communicable Diseases in Europe. ECDC, Stockholm, Sweden.
- Furet, J.P., Firmesse, O., Gourmelon, M., Bridonneau, C., Tap, J., Mondot, S., Dore, J. and Corthier, G., 2009. Comparative assessment of human and farm animal faecal microbiota using real-time quantitative PCR. *FEMS Microbiology Ecology* 68: 351-362.
- Guarner, F., Sanders, M.E., Eliakim, R., Fedorak, R., Gangl, A., Garisch, J., Kaufmann, P., Karakan, T., Khan, A.G. and Kim, N., 2017. Probiotics and prebiotics. World Gastroenterology Organisation Global Guidelines. World Gastroenterology Organisation, Milwaukee, WI, USA. Available at: <https://tinyurl.com/4sapeppc>
- Guo, X., Xia, X., Tang, R., Zhou, J., Zhao, H. and Wang, K., 2008. Development of a real-time PCR method for *Firmicutes* and *Bacteroidetes* in faeces and its application to quantify intestinal population of obese and lean pigs. *Letters in Applied Microbiology* 47: 367-373.
- Hartemink, R., Domenech, V.R. and Rombouts, F.M., 1997. LAMVAB – A new selective medium for the isolation of lactobacilli from faeces. *Journal of Microbiological Methods* 29: 77-84.
- Hill, C., Guarner, F., Reid, G., Gibson, G.R., Merenstein, D.J., Pot, B., Morelli, L., Canani, R.B., Flint, H.J., Salminen, S., Calder, P.C. and Sanders, M.E., 2014. Expert consensus document. The International Scientific Association for Probiotics and Prebiotics consensus statement on the scope and appropriate use of the term probiotic. *Nature Reviews Gastroenterology and Hepatology* 11: 506-514.
- Kalliomaki, M., Salminen, S., Arvilommi, H., Kero, P., Koskinen, P. and Isolauri, E., 2001. Probiotics in primary prevention of atopic disease: a randomised placebo-controlled trial. *The Lancet* 357: 1076-1079.
- Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M. and Glockner, F.O., 2013. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Research* 41: e1.
- Kotowska, M., Albrecht, P. and Szajewska, H., 2005. *Saccharomyces boulardii* in the prevention of antibiotic-associated diarrhoea in children: a randomized double-blind placebo-controlled trial. *Alimentary Pharmacology and Therapeutics* 21: 583-590.
- Kozich, J.J., Westcott, S.L., Baxter, N.T., Highlander, S.K. and Schloss, P.D., 2013. Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform. *Applied and Environmental Microbiology* 79: 5112-5120.
- Ladirat, S.E., Schoterman, M.H., Rahaoui, H., Mars, M., Schuren, F.H., Gruppen, H., Nauta, A. and Schols, H.A., 2014a. Exploring the effects of galacto-oligosaccharides on the gut microbiota of healthy adults receiving amoxicillin treatment. *British Journal of Nutrition* 112: 536-546.
- Ladirat, S.E., Schuren, F.H., Schoterman, M.H., Nauta, A., Gruppen, H. and Schols, H.A., 2014b. Impact of galacto-oligosaccharides on the gut microbiota composition and metabolic activity upon antibiotic treatment during *in vitro* fermentation. *FEMS Microbiology Ecology* 87: 41-51.

- Lazo-Vélez, M.A., Serna-Saldívar, S.O., Rosales-Medina, M.F., Tinoco-Alvear, M. and Briones-García, M., 2018. Application of *Saccharomyces cerevisiae* var. *boulardii* in food processing: a review. *Journal of Applied Microbiology* 125: 943-951.
- Legen, I., Kracun, M., Salobir, M. and Kerc, J., 2006. The evaluation of some pharmaceutically acceptable excipients as permeation enhancers for amoxicillin. *International Journal of Pharmaceutics* 308: 84-89.
- Liu, L., Firman, J., Tanes, C., Bittinger, K., Thomas-Gahring, A., Wu, G.D., Van den Abbeele, P. and Tomasula, P.M., 2018. Establishing a mucosal gut microbial community in vitro using an artificial simulator. *PLoS ONE* 13: e0197692.
- Mantegazza, C., Molinari, P., D'Auria, E., Sonnino, M., Morelli, L. and Zuccotti, G.V., 2018. Probiotics and antibiotic-associated diarrhea in children: a review and new evidence on *Lactobacillus rhamnosus* GG during and after antibiotic treatment. *Pharmacological Research* 128: 63-72.
- McFarland, L.V., 2008. Antibiotic-associated diarrhea: epidemiology, trends and treatment. *Future Microbiology* 3: 563-578.
- McFarland, L.V. and Elmer, G.W., 2005. Properties of evidence-based probiotics for human health. In: Goktepe, I., Juneja, V.K. and Ahmedna, M. (eds.) *Probiotics in food safety and human health*. CRC Press, Boca Raton, FL, USA, pp. 110-128.
- McFarland, L.V., Surawicz, C.M., Greenberg, R.N., Elmer, G.W., Moyer, K.A., Melcher, S.A., Bowen, K.E. and Cox, J.L., 1995. Prevention of beta-lactam-associated diarrhea by *Saccharomyces boulardii* compared with placebo. *American Journal of Gastroenterology* 90: 439-448.
- Minekus, M., Alming, M., Alvito, P., Ballance, S., Bohn, T., Bourlieu, C., Carrière, F., Boutrou, R., Corredig, M., Dupont, D., Dufour, C., Egger, L., Golding, M., Karakaya, S., Kirkhus, B., Le Feunteun, S., Lesmes, U., Macierzanka, A., Mackie, A., Marze, S., McClements, D.J., Ménard, O., Recio, I., Santos, C.N., Singh, R.P., Vegarud, G.E., Wickham, M.S., Weitschies, W. and Brodkorb, A., 2014. A standardised static *in vitro* digestion method suitable for food – an international consensus. *Food and Function* 5: 1113-1124.
- Minekus, M., Smeets-Peters, M., Bernalier, A., Marol-Bonnin, S., Havenaar, R., Marteau, P., Alric, M., Fonty, G. and Huis in't Veld, J.H., 1999. A computer-controlled system to simulate conditions of the large intestine with peristaltic mixing, water absorption and absorption of fermentation products. *Applied Microbiology and Biotechnology* 53: 108-114.
- Moens, F., Duysburgh, C., van den Abbeele, P., Morera, M. and Marzorati, M., 2019. *Lactobacillus rhamnosus* GG and *Saccharomyces cerevisiae* *boulardii* exert synergistic antipathogenic activity *in vitro* against enterotoxigenic *Escherichia coli*. *Beneficial Microbes* 10: 923-935.
- Molly, K., Vande Woestyne, M. and Verstraete, W., 1993. Development of a 5-step multi-chamber reactor as a simulation of the human intestinal microbial ecosystem. *Applied Microbiology and Biotechnology* 39: 254-258.
- Moré, M.I. and Swidsinski, A., 2015. *Saccharomyces boulardii* CNCM I-745 supports regeneration of the intestinal microbiota after diarrheic dysbiosis – a review. *Clinical and Experimental Gastroenterology* 8: 237-255.
- Neut, C., Mahieux, S. and Dubreuil, L.J., 2017. Antibiotic susceptibility of probiotic strains: Is it reasonable to combine probiotics with antibiotics? *Médecine et Maladies Infectieuses* 47: 477-483.
- Palleja, A., Mikkelsen, K., Forslund, S., Kashani, A., Allin, K., Nielsen, T., Hansen, T., Feng, Q., Zhang, C., Pyl, P., Coelho, L.P., Yang, H., Wang, J., Typas, A., Frost, M., Nielsen, H., Bork, P., Wang, J. and Pedersen, O., 2018. Recovery of gut microbiota of healthy adults following antibiotic exposure. *Nature Microbiology* 3: 1255-1265.
- Panda, S., El khader, I., Casellas, F., López Vivancos, J., García Cors, M., Santiago, A., Cuenca, S., Guarner, F. and Manichanh, C., 2014. Short-term effect of antibiotics on human gut microbiota. *PLoS ONE* 9: e95476.
- Possemiers, S., Verthe, K., Uyttendaele, S. and Verstraete, W., 2004. PCR-DGGE-based quantification of stability of the microbial community in a simulator of the human intestinal microbial ecosystem. *FEMS Microbiology Ecology* 49: 495-507.
- Pozzoni, P., Riva, A., Bellatorre, A.G., Amigoni, M., Redaelli, E., Ronchetti, A., Stefani, M., Tironi, R., Molteni, E.E., Conte, D., Casazza, G. and Colli, A., 2012. *Saccharomyces boulardii* for the prevention of antibiotic-associated diarrhea in adult hospitalized patients: a single-center, randomized, double-blind, placebo-controlled trial. *American Journal of Gastroenterology* 107: 922-931.
- Ramamurthy, D., Pazhani, G.P., Sarkar, A., Nandy, R.K., Rajendran, K., Sur, D., Manna, B. and Ramamurthy, T., 2013. Case-control study on the role of enterotoxigenic *Bacteroides fragilis* as a cause of diarrhea among children in Kolkata, India. *PLoS ONE* 8: e60622.
- Rinttilä, T., Kassinen, A., Malinen, E., Krogus, L. and Palva, A., 2004. Development of an extensive set of 16S rDNA-targeted primers for quantification of pathogenic and indigenous bacteria in faecal samples by real-time PCR. *Journal of Applied Microbiology* 97: 1166-1177.
- Schloss, P.D. and Westcott, S.L., 2011. Assessing and improving methods used in operational taxonomic unit-based approaches for 16S rRNA gene sequence analysis. *Applied and Environmental Microbiology* 77: 3219-3226.
- Soldi, S., Vasileiadis, S., Lohner, S., Uggeri, F., Puglisi, E., Molinari, P., Donner, E., Sieland, C., Decsi, T., Sailer, M. and Theis, S., 2019. Prebiotic supplementation over a cold season and during antibiotic treatment specifically modulates the gut microbiota composition of 3-6 year-old children. *Beneficial Microbes* 10: 253-263.
- Sommer, F. and Bäckhed, F., 2013. The gut microbiota – masters of host development and physiology. *Nature Reviews Microbiology* 11: 227-238.
- Spacova, I., O'Neill, C. and Lebeer, S., 2020. *Lactocaseibacillus rhamnosus* GG inhibits infection of human keratinocytes by *Staphylococcus aureus* through mechanisms involving cell surface molecules and pH reduction. *Beneficial Microbes* 11: 703-715.
- Szajewska, H. and Kolodziej, M., 2015. Systematic review with meta-analysis: *Lactobacillus rhamnosus* GG in the prevention of antibiotic-associated diarrhoea in children and adults. *Alimentary Pharmacology and Therapeutics* 42: 1149-1157.

- Tremblay, A., Fatani, A., Ford, A.L., Piano, A., Nagulesapillai, V., Auger, J., MacPherson, C.W., Christman, M.C., Tompkins, T.A. and Dahl, W.J., 2020. Safety and effect of a low- and high-dose multi-strain probiotic supplement on microbiota in a general adult population: a randomized, double-blind, placebo-controlled study. *Journal of Dietary Suppl.* 18: 227-247.
- Tsitko, I., Wiik-Miettinen, F., Mattila, O., Rosa-Sibakov, N., Seppänen-Laakso, T., Maukonen, J., Nordlund, E. and Saarela, M., 2019. A small *in vitro* fermentation model for screening the gut microbiota effects of different fiber preparations. *International Journal of Molecular Sciences* 20: 1925.
- Van de Wiele, T., Van den Abbeele, P., Ossieur, W., Possemiers, S. and Marzorati, M., 2015. The simulator of the human intestinal microbial ecosystem (SHIME®). In: Verhoeckx, K., Cotter, P., López-Expósito, I., Kleiveland, C., Lea, T., Mackie, A., Requena, T., Swiatecka D. and Wichers, H. (eds.), *The impact of food bioactives on health: in vitro and ex vivo models*. Springer International Publishing, Berlin, Germany, pp. 305-317.
- Van den Abbeele, P., Belzer, C., Goossens, M., Kleerebezem, M., De Vos, W.M., Thas, O., De Weirtdt, R., Kerckhof, F.M. and Van de Wiele, T., 2013. Butyrate-producing *Clostridium* cluster XIVa species specifically colonize mucins in an *in vitro* gut model. *ISME Journal* 7: 949-961.
- Van den Abbeele, P., Grootaert, C., Marzorati, M., Possemiers, S., Verstraete, W., Gerard, P., Rabot, S., Bruneau, A., El Aidy, S., Derrien, M., Zoetendal, E., Kleerebezem, M., Smidt, H. and Van de Wiele, T., 2010. Microbial community development in a dynamic gut model is reproducible, colon region specific, and selective for *Bacteroidetes* and *Clostridium* cluster IX. *Applied and Environmental Microbiology* 76: 5237-5246.
- Van den Abbeele, P., Roos, S., Eeckhaut, V., MacKenzie, D.A., Derde, M., Verstraete, W., Marzorati, M., Possemiers, S., Vanhoecke, B., Van Immerseel, F. and Van de Wiele, T., 2012. Incorporating a mucosal environment in a dynamic gut model results in a more representative colonization by lactobacilli. *Microbial Biotechnology* 5: 106-115.
- Van den Abbeele, P., Taminiau, B., Pinheiro, I., Duysburgh, C., Jacobs, H., Pijls, L. and Marzorati, M., 2018. Arabinosylo-oligosaccharides and inulin impact inter-individual variation on microbial metabolism and composition, which immunomodulates human cells. *Journal of Agricultural and Food Chemistry* 66: 1121-1130.
- Van Herreweghen, F., Van den Abbeele, P., De Mulder, T., De Weirtdt, R., Geirnaert, A., Hernandez-Sanabria, E., Vilchez-Vargas, R., Jauregui, R., Pieper, D.H., Belzer, C., De Vos, W.M. and Van de Wiele, T., 2017. *In vitro* colonisation of the distal colon by *Akkermansia muciniphila* is largely mucin and pH dependent. *Beneficial Microbes* 8: 81-96.
- Vanderhoof, J.A., Whitney, D.B., Antonson, D.L., Hanner, T.L., Lupo, J.V. and Young, R.J., 1999. *Lactobacillus* GG in the prevention of antibiotic-associated diarrhea in children. *Journal of Pediatrics* 135: 564-568.
- Wang, Q., Garrity, G.M., Tiedje, J.M. and Cole, J.R., 2007. Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Applied and Environmental Microbiology* 73: 5261-5267.
- White, A.R., Kaye, C., Poupard, J., Pypstra, R., Woodnutt, G. and Wynne, B., 2004. Augmentin® (amoxicillin/clavulanate) in the treatment of community-acquired respiratory tract infection: a review of the continuing development of an innovative antimicrobial agent. *Journal of Antimicrobial Chemotherapy* 53: i3-i20.